# Effect of dietary organic trace minerals for sows on piglet vitality and milk composition

CPH Pig seminar 2<sup>nd</sup> of February 2023

#### Anja Varmløse Strathe

Associate Professor

Department of Veterinary and Animal Sciences

Section for Production, Nutrition and Health

avst@sund.ku.dk

UNIVERSITY OF COPENHAGEN



The project is a collaboration between SEGES and KU Funded by:

Norma and Frode S. Jacobsens Fund

## Are there differences between different types of organic trace minerals?

Higher availability in organic vs. inorganic trace minerals (TM)

- Many types of organic TM -> many price levels
  - Glycinated (cheaper type): Mineral bound in complex with glycine
  - Amino acid chelate (more expensive): Mineral is bound to amino acid

In this study we focus on cupper, manganese and zinc

# Why are cupper, zinc and manganese important to the sow?

Important for claw health -> improved longevity of sows?

Higher availability of Zn, Mn and Cu:

- Fewer piglets with low birth weight, higher litter size and weight at farrowing, more vital piglets
  - Many TM are transported from sow to foetuses and colostrum
- Impact on milk production -> higher weaning weight -> better development? and more prepared for weaning?
  - The transfer of TM to milk is more limited than for colostrum

## **Experimental design**

Sows were fed experimental diets from previous farrowing -> next weaning

- Two groups:
  - Organic glycinated TM
  - Organic amino acid chelated TM

Standard diet	Glycinated TM diet	Amino acid chelated TM diet
Copper sulphate	10 mg/kg copper chelate glycine hydrate	10 mg/kg copper amino acid chelate
Manganese oxide	20 mg/kg Manganese chelate glycine hydrate	20 mg/kg manganese amino acid chelate
Zinc oxide	50 mg/kg zinc glycine chelate	50 mg/kg zinc amino acid chelate

## Effect on piglet vitality at farrowing

 743 piglets from 39 litters were weighed and measured individually within 8 hours of birth

- Measures of piglet vitality:
  - BW at birth -> not the best predictor of performance
  - BMI = birth weight / crown-rump length<sup>2</sup> -> good predictor of performance
  - Abdominal circumference -> good predictor of performance
  - Ponderal Index = birth weight / crown-rump length<sup>3</sup>
  - IUGR-score -> not very objective
  - Crown-rump length

#### More results will come...

- SEGES Innovation:
  - Claw health before and after use of different organic TM
  - Birth weight of piglets
  - ADG of the litter, litter size and litter weight at weaning



# Thank you for listening!

The project was a collaboration between SEGES Innovation and KU

#### The project was funded by:

Norma and Frode S. Jacobsens Fund



